

ALEXSANDER FERRAZ

*Universidade Federal de Pelotas, UFPEL, Pelotas, RS,
Brasil.*

RENATA FONTES ONGARATTO

*Universidade Federal de Pelotas, UFPEL, Pelotas, RS,
Brasil.*

CLEBER MARTINS RIBEIRO

*Universidade Federal de Pelotas, UFPEL, Pelotas, RS,
Brasil.*

TIAGO FELIPE BARBOSA MOREIRA

*Universidade Federal de Pelotas, UFPEL, Pelotas, RS,
Brasil.*

RODRIGO LEITE DOS SANTOS

*Universidade Federal de Pelotas, UFPEL, Pelotas, RS,
Brasil.*

WESLEY OLIVEIRA

*Universidade Federal de Pelotas, UFPEL, Pelotas, RS,
Brasil.*

GABRIELA DE ALMEIDA CAPELLA

*Universidade Federal de Pelotas, UFPEL, Pelotas, RS,
Brasil.*

MARIANA TIMM KROLOW

*Universidade Federal de Pelotas, UFPEL, Pelotas, RS,
Brasil.*

THAÍS REICHOW

*Universidade Federal de Pelotas, UFPEL, Pelotas, RS,
Brasil.*

LEANDRO QUINTANA NIZOLI

*Universidade Federal de Pelotas, UFPEL, Pelotas, RS,
Brasil.*

*Recebido em março de 2024.
Aprovado em março de 2024.*

POTENTIAL ZOOONOTIC PARASITES OF DOGS FECES IN THE VICINITY OF SCHOOLS IN THE CITY OF CAPÃO DO LEÃO, RIO GRANDE DO SUL, BRAZIL

ABSTRACT

The aim of this work was to identify the presence of evolutive forms of parasites with zoonotic potential, in dog's feces, found in the vicinity of urban schools in the city of Capão do Leão, in Rio Grande do Sul. Samples have been collected from seven schools and analyzed using the coproparasitological techniques of Willis Mollay, Faust and Hoffmann, Pns and Janer. Of the analyzed samples, 83.3% (25/30) were positive for some genus of gastrointestinal parasite. *Ancylostoma* spp. was the prevalent genus, being observed in 63.3% of the samples. There were still observed eggs of *Trichuris vulpis* (16.7%), *Toxocara* spp. (6.7%), *Spirometra* sp. (6.7%), *Dipylidium caninum* (3.3%), *Giardia* sp. (10%) and *Cystoisospora* spp. (6.7%). These results highlight the need for practices that prevent and control parasitic infections, followed by the responsible pet ownership and control of semi-domiciled or abandoned animals, in order to reduce the spread of these agents in the environment, once the presence of evolutive forms of parasites with potential zoonotic characterizes a risk to the population and other animals.

Keywords: environmental contamination. zoonoses. companion animals. parasites.

PARASITOS POTENCIALMENTE ZOOONÓTICOS EM FEZES DE CÃES NAS PROXIMIDADES DE ESCOLAS DO MUNICÍPIO DE CAPÃO DO LEÃO, RIO GRANDE DO SUL, BRASIL

RESUMO

O objetivo deste trabalho foi identificar a presença de formas evolutivas de parasitos com potencial zoonótico, em fezes de cães, encontradas nas proximidades de escolas urbanas do município de Capão do Leão, no Rio Grande do Sul. As amostras foram coletadas em sete escolas e analisadas através das técnicas coproparasitológicas de Willis Mollay, Faust e Hoffmann, Pons e Janer. Das amostras analisadas, 83,3% (25/30) foram positivas para algum gênero de parasito gastrintestinal. *Ancylostoma* spp. foi o gênero prevalente, sendo observado em 63,3% das amostras. Ainda foram observados, ovos de *Trichuris vulpis* (16,7%), *Toxocara* spp. (6,7%), *Spirometra* spp. (6,7%), *Dipylidium caninum* (3,3%), cistos de *Giardia* sp. (10%) e oocistos de *Cystoisospora* spp. (6,7%). Esses resultados evidenciam a necessidade de práticas que previnam e controlem as infecções parasitárias, acompanhado da guarda responsável dos animais domiciliados e controle dos semidomiciliados ou abandonados, de modo que diminuam a propagação destes agentes no ambiente, pois a presença de formas evolutivas de parasitos com potencial zoonótico caracteriza um risco para a população e outros animais.

Palavras-Chave: contaminação ambiental. zoonoses. animais de companhia. parasitos.

INTRODUÇÃO

The human and animal relationship, mainly with cats and dogs, promotes countless benefits of psychological, physiological and sociological for the tutors. It is estimated that 46.1% of the Brazilian houses possess at least one dog and 19.3% have cats (IBGE, 2020). However, infected animals are responsible for parasitic zoonoses dissemination, mostly through soil fecal contamination, representing a risk to human health (LIMA et al., 2017). As a consequence, it creates an important public health problem, correlated with economical, sociocultural and environmental factors (CASSENOTE et al., 2011).

The most affected individuals by parasitic diseases are scholar and pre scholar children, due to countless factors such as: more contact with dirt and sand; walking shoeless and geophagy habits; precarity on maintaining personal hygiene practices and immune system immaturity (CAVAGNOLLI et al., 2015). The occurrence of parasitic diseases on infant age can create nutritional deficiency nutrition, leading to gastrointestinal disorders, with the presence of chronic diarrhea. So, it directly affects the children's physical and intellectual development besides interfering on school yield (NAVONE et al., 2017; PEDRAZA, 2014).

Dogs and cats perform an important role as definitive host of countless parasites, eliminating helminth eggs, cysts and oocysts of protozoans through feces, leading to environmental contamination and aiding zoonosis dissemination. Among the factors that collaborate for soil contamination stands out the presence of semi-domiciled and walking dogs and cats in urban areas and the absence of the tutor's habits on collecting the dejects of their animals (CASSENOTE et al., 2011).

It is evident that infected dogs and cats represent a relevant problem of public health, due the possibility of zoonoses transmission, as cutaneous *larva migrans* (CLM) and visceral *larva migrans* (VLM), caused by helminths *Ancylostoma* spp. and *Toxocara* spp., respectively (CASSENOTE et al., 2011; SANTOS et al., 2009); besides the giardiasis infirmity caused by the zoonotic protozoan *Giardia* spp. (SAVIOLI, 2006).

It is relevant to highlight, the cutaneous *larva migrans*, also known as "geographic worm", it is caused by the penetration of infective larvae of *Ancylostoma* spp. on the epithelial layer of human skin, triggering a self-limited cutaneous inflammation, with serpiginous, erythematous and pruritic injuries on inferior limbs, buttocks and hands, mainly (SANTARÉM, 2004). As for the visceral *larva migrans* occurs by the migration of *Toxocara* spp. larvae on the tissues of non-habitual hosts, as human, and triggers inflammatory injuries on viscera, in which can cause damage on important organs as liver, brain and eyes (SANTOS et al., 2009).

In that way, the aim of this work was to identify the presence of evolutive forms of parasites with zoonotic potential, in dog's feces, found in the vicinity of urban schools at the city of Capão do Leão, in Rio Grande do Sul, Brazil.

MATERIALS AND METHODS

Feces samples have been collected in the vicinity of seven urban schools in the city of Capão do Leão (Latitude: 31°45'48" S, Longitude: 52°29'02" W), located on the South region of the state of Rio Grande do Sul. The city has an estimated population of 25.462 people (IBGE, 2021). All of the samples collected were fresh and were packed in individual plastic bags, identified and stored in isothermal boxes for further analysis at the Laboratory of Parasites Diseases (LADOPAR) at the Veterinary College of Federal University of Pelotas (UFPe1).

The coproparasitological techniques used for the analysis of the samples were: Willis-Mollay (1921), which consists on the flotation of light eggs of helminths on supersaturated solution; Faust (1938), a centrifuge technique on 33% zinc sulfate

solution, used to search for cysts and oocysts of *Giardia* sp. and the spontaneous sedimentation technique of Hoffmann, Pons and Janer (1934), for the research of heavy eggs, as cestodes and trematodes. The genera differentiation of enteroparasites, was given through the morphological characteristics of eggs, cysts and oocysts, visualized on optical microscopy, on 100x and 400x magnification.

RESULTS

Of the 30 samples analyzed, 83.3% (25/30) were positive for at least one genus of enteroparasites. Of those, 16 (64%) presented monoparasitism and 9 (36%) contained association of parasites. *Ancylostoma* spp. eggs were the most prevalent, being observed in 63.3% (19/30) of the samples. There were still found eggs of *Toxocara* spp., *Trichuris vulpis*, *Dipylidium caninum* and *Spirometra* sp., oocysts of *Cystoisospora* spp. and cysts of *Giardia* sp. The association of *Ancylostoma* spp. + *Giardia* sp. was the most observed (10%) (Table 1).

Table 1. Prevalence of parasites found on dog's fecal samples in the vicinity of urban schools in the city of Capão do Leão (RS).

Parasite	Positive samples n(%)
Monoparasitism	
<i>Ancylostoma</i> spp.	11(36.6)
<i>Trichuris vulpis</i>	2(6.7)
<i>Cystoisospora</i> spp.	2(6.7)
<i>Toxocara</i> spp.	1(3.3)
Multiparasitism	
<i>Ancylostoma</i> spp. + <i>Giardia</i> sp.	3(10.0)
<i>Ancylostoma</i> spp. + <i>Trichuris vulpis</i>	2(6.7)
<i>Ancylostoma</i> spp. + <i>Toxocara</i> spp.	1(3.3)
<i>Ancylostoma</i> spp. + <i>Spirometra</i> sp.	1(3.3)
<i>Trichuris vulpis</i> + <i>Spirometra</i> sp.	1(3.3)
<i>Ancylostoma</i> spp. + <i>Dipylidium caninum</i>	1(3.3)
Total	25(83,3)

DISCUSSION

Dogs can host several genera of enteric parasites and most of them present transmission potential to humans, representing a public health problem (TRAVERSA, 2014). In that way, several studies are being carried out to determinate the prevalence of zoonotic parasites on dog's fecal samples collected in public places, such as Bricarello et al. (2018) in beaches of Florianópolis (SC), Providelo et al. (2020) in public parks of Botucatu (SP) and Ribeiro et al. (2013) in parks of the city of Esteio (RS).

Ancylostoma was the most prevalent genus, being observed in 63.3% of the samples. Other authors also report the predominance of this parasite on their studies, like Campos Filho (2008) at Bahia (47.9%), Bricarello et al. (2020) in Santa Catarina (42.2%) and Coronato et al. (2012) in Rio de Janeiro (91.9%). *Ancylostoma* spp., is a nematode of great importance in public health, because it is the agent responsible for the parasitic zoonosis named cutaneous larva *migrans*, commonly known as "geographic worm", that occurs by skin penetration of third-stage larvae (infective form) of the parasite. After the penetration on the epidermis, the larvae migrate on the subcutaneous tissue, promoting serpiginous eruptions, mostly on the locations with greater contact

with the contaminated mean, such as feet, legs, buttocks and hand (ALMEIDA et al., 2007). Araújo et al. (2000), reported the occurrence of cutaneous *larva migrans* in children of an infant education school at Campo Grande, MS, where of the 16 students that attended the school, 37.5% (6/16) acquired this parasite infirmity through the contact with contaminated sand by the cat's feces on recreational areas.

Toxocara spp. eggs were found in 6.7% of the samples. The presence of the eggs of this parasite indicates human visceral *larva migrans* and ocular *larva migrans* syndrome infection, which occurs by the ingestion of eggs containing the infective larvae of the parasite and further migration on different organs (OVERGAAUW and VAN KNAPEN, 2013). Scaini et al. (2003) reported the presence of this parasite in 9.3% of the dogs fecal samples collected in Balneario Cassino, Rio Grande do Sul, a percentage slightly higher than our study. An inferior prevalence was observed by Bricarello et al. (2018) in Florianópolis (SC) e Matesco et al. (2006) in Porto Alegre (RS) that found a positivity of 0.9% and 0.8% respectively for this genus of parasite.

Cysts of *Giardia* sp. were found in 10% of the samples. This protozoan also possesses a zoonotic potential, being a source of infection for human beings and other animals. (SILVA and ARAÚJO, 2013). The infection occurs from the ingestion of infectant cysts present on contaminated water and food. The World Organization of Health, estimates that around 280 million of people around the world are stricken each year by this protozoan, mostly children (FRASSON et al., 2010). In Ribeirão Preto/SP, Zagui et al. (2017) on a study of intestinal parasite diseases in children of a philanthropic institution diagnosed that 21.2% (14/66) of the children were infected by *Giardia* spp.

Trichuris vulpis was observed in 16.7% of the samples. Alves et al. (2014) also reported the presence of *T. vulpis* eggs in fecal samples from public parks of Pindamonhangaba (SP) and Leite et al. (2013) in public ways of Guarapuava (PR), that found, respectively 3.1% e 11.9% of positive samples. This nematode parasitizes the large intestine of dogs, causing, mostly on animals with high parasite load, anemia, diarrhea, bloody diarrhea, vomit, pain, abdominal distention and some dogs can present rectal prolapse (SHERDING and JOHNSON, 2008). The infection occurs through the ingestion of eggs containing the infectant larvae present on the soil or contaminated food with the infected animals feces (DI CESARE et al., 2012). Although it is low described, some works report the infection by *T. vulpis* in humans, indicating its zoonotic potential, as Márquez-Navarro et al., (2012) that reported a case in a kid in Mexico and Dunn et al. (2002) that reported a case of *T. vulpis* in a woman with duodenal ulcer and chronic diarrhea.

Dipylidium caninum, a cestode that has fleas and lice as intermediate hosts, was found in only one sample (3.3%). Low percentages were also evidenced by Rosales and Malheiros (2017) in Pantanal and Junior et al. (2015) at the city of Natal (RN), where both found 1.7% of positive samples. The low prevalence of *D. caninum* can be justified by the fact that the ovigerous capsule of the parasite are hardly found during the coproparasitological exam, and the best way to diagnose is through the visualization of the proglottids on fresh feces or the founding of adult forms on necropsies (GENNARI, 1999). Some reports describe human infections by *D. caninum*, mainly in children (JIANG et al., 2017; NARASIMHAM et al., 2013).

Spirometra sp. eggs and *Cystoisospora* spp. oocysts were present in 6.7% of each sample. Other authors also reported the presence of these parasites on their works (FERRAZ et al., 2021, ROSALES and MALHEIROS, 2017). *Spirometra* sp., is a cestode that attacks the small intestine of several mammals, as dogs and cats, being able to also parasite humans. It is the responsible for the parasitic zoonosis named Sparganosis and infection occurs through ingestion of the infected intermediate host. (TANTALEAN et al., 2005). As for the *Cystoisospora* spp., although does not present zoonotic potential,

can cause intestinal injuries on the host, leading to diarrhea and impaired nutrient absorption cases, especially in puppies (BARUTZKI et al., 2013).

The adoption of sanitary measures, besides being beneficial to animal health, also has a great importance for public health, as there will be a reduction of spread of zoonotic parasites. On these way, it is necessary to emphasize the importance of sanitary care and the responsible pet ownership, once the unfamiliarity of the population on the risks involved it is one of the mainly factors that trigger these diseases (PERUCA et al., 2009). Researches show that the majority of Brazilian citizens are unfamiliar on the risks to health related to dog's feces left on public soils (PEREIRA et al., 2016; SIMONATO et al., 2019).

CONCLUSIONS

It was observed the high prevalence of the parasites with zoonotic potential in the vicinity of schools of the city of Capão do Leão. These results highlighted the need to adopt preventive sanitary measures, such as responsible pet ownership, as well the control of semi-domiciled and abandoned cats and dogs, always focusing on the well-being and quality of life of all animals.

It is also extremely important to carry out scientific studies that aim to document the risks involving parasitic diseases in the most diverse urban locations, avoiding the spread of those.

REFERÊNCIAS

- ALMEIDA, A.B.P.F. et al. Contaminação por fezes caninas das praças públicas de Cuiabá, Mato Grosso. *Brazilian Journal Veterinary Research and Animal Science*, v. 44, n. 2, p.132-136, 2007.
- ARAÚJO, F.R. et al. Larva migrans cutânea em crianças de uma escola em área do Centro-Oeste do Brasil. *Revista de Saúde Pública*, v.34, n.1, p.84-85, 2000. <Doi: 10.1590/S0034-89102000000100015>
- BARUTZKI, D.; SCHAPER, R. Prevalence of Endoparasites in Young Dogs and Cats up to One Year of Age. *Parasitology Research*, v.112, n.1, p.119-131, 2013. <Doi: 10.1007/s00436-013-3286-6>
- BRICARELLO P.A. et al. Potential zoonotic parasites in dog and cat feces from three beaches and surrounding areas of Greater Florianópolis, Santa Catarina. *Brazilian Journal of Veterinary Research and Animal Science*, v.57, n.3, p.1-9, 2020. <DOI: 10.11606/issn.1678-4456.bjvras.2020.168442>.
- BRICARELLO, P.A. et al. Contamination by parasites of zoonotic importance in fecal samples from Florianópolis Beaches, Santa Catarina State. Brazil. *Brazilian Journal of Veterinary Research and Animal Science*, v.55, n.1, p.1-10, 2018. <DOI: 10.11606/issn.1678-4456.bjvras.2018.133259>.
- CAMPOS FILHO, P.C. et al. Parasitas zoonóticos em fezes de cães em praças públicas do município de Itabuna, Bahia, Brasil. *Revista Brasileira de Parasitologia Veterinária*, v.17, n.4, p.206-209, 2008. <DOI: 10.1590/S1984-29612008000400007>.
- CASSENOTE, A.J.F. et al. Contaminação do solo por ovos de geo-helminthos com potencial zoonótico na municipalidade de Fernandópolis, Estado de São Paulo, entre 2007 e 2008. *Revista da Sociedade Brasileira Medicina Tropical*, v.44, n.3, p.371-374, 2011. <DOI: 10.1590/S0037-86822011005000026>.

POTENTIAL ZOOONOTIC PARASITES OF DOGS FECES IN THE VICINITY OF SCHOOLS IN THE CITY OF CAPÃO DO LEÃO, RIO GRANDE DO SUL, BRAZIL / PARASITOS POTENCIALMENTE ZOOONÓTICOS EM FEZES DE CÃES NAS PROXIMIDADES DE ESCOLAS DO MUNICÍPIO DE CAPÃO DO LEÃO, RIO GRANDE DO SUL, BRASIL

- CASTRO, C.; OLIVEIRA, J.B.; HERNÁNDEZ, J. et al. Contaminación por parásitos gastrointestinales de caninos en dieciocho playas del Pacífico Central de Costa Rica: implicaciones para la salud pública. *Revista Brasileira de Ciência Veterinária*, v.27, n.2, p.7-16, 2009.
- CAVAGNOLLI, N.I.; CAMELLO, J.T.; TESSER, S. et al. Prevalência de enteroparasitoses e análise socioeconômica de escolares em Flores da Cunha-RS. *Revista de Patologia Tropical*, v. 44, n.3, p. 312-322, 2015. <DOI: 10.5216/rpt.v44i3.38018>.
- CORONATO, B.; BASTOS, O.M.P.; DUARTE, R. et al. Parasites in stool samples in the environment of Ilha da Marambaia, Rio de Janeiro, Brazil: an approach in public health. *Revista do Instituto de Medicina Tropical de São Paulo*, v. 54, n. 2, p. 65-68, 2012. <DOI: 10.1590/S0036-46652012000200002>.
- DI CESARE, A. et al. Mixed trichuroid infestation in a dog from Italy. *Parasites & Vectors*, v.5, p.1-6, 2012. <DOI: 10.1186 /1756-3305-5-128>.
- DIBA, V.C.; WHITTY, C.J.M.; GREEN, T. Cutaneous larva migrans acquired in Britain. *Clinical and experimental dermatology*, v.29, n.5, p.555-556, 2004.
- DUNN, J.J. et al. *Trichuris vulpis* recovered from a patient with chronic diarrhea and five dogs. *Journal of Clinical Microbiology*, v.40, n.7, p.2703-4, 2002. <DOI: 10.1128/JCM.40.7.2703-2704.2002>.
- FAUST, E.C.; D'ANTONI, J.S.; ODOM, V. A critical study of clinical laboratory technics for the diagnosis of protozoan cysts and helminth eggs in feces I. Preliminary communication. *American Journal of Tropical Medicine*, v18, n.2, p.169-183, 1938.
- FERRAZ, A. et al. Prevalência de parasitos potencialmente zoonóticos nas proximidades das escolas urbanas do município de Pedro Osório, Rio Grande do Sul, Brasil. *Revista UNILUS Ensino e Pesquisa*, v.18, n.51, p.77-84, 2021.
- FRASSON, A.P. et al. *Giardia lamblia*: Distribuição de microtúbulos no citoesqueleto de trofozoítos e cistos utilizando taxóide fluorescente. *Revista de Patologia Tropical*, v.39, n.1, p.21-32, 2010. >DOI: doi.org/10.5216/rpt.v39i1.9495>.
- GENNARI, S.M. et al. Occurrence of protozoa and helminths in faecal samples of dogs and cats from São Paulo city. *Brazilian Journal of Veterinary Research and Animal Science*, v.36, n.2, p.87-91, 1999. <DOI: 10.1590/S1413-95961999000200006>.
- HOFFMAN, W.A.; PONS, J.A.; JANER, J.L. Sedimentation concentration method in *Schistosomiasis mansoni*. *The Puerto Rico Journal of Public Health and Tropical Medicine*, v.9, p.283-298, 1934.
- INSTITUTO BRASILEIRO DE GEOGRAFIA E ESTATISTICA (IBGE). Disponível em: <https://www.ibge.gov.br/cidades-e-estados/rs/capao-do-leao.html>. Acesso em 02 mai. 2021.
- JIANG, P. et al. A Human Case of Zoonotic Dog Tapeworm, *Dipylidium caninum* (Eucestoda: Dilepidiidae), in China. *The Korean Journal of Parasitology*, v.55, n.1, p.61-64, 2017. <DOI: 10.3347/kjp.2017.55.1.61>.
- JUNIOR, A.L.F.A.; ARAÚJO, K.B.S.; MEDEIROS, V.S. Ocorrência de parasitas com potencial zoonótico em fezes de cães coletadas em vias públicas da cidade de Natal. *Revista Humano Ser*, v.1, n.1, p. 52-59, 2015.
- LIMA, F.S.C. et al. Contaminação ambiental por ovos de *Ancylostoma* spp. e *Toxocara* spp. em áreas de seis praças públicas do município de Valença, estado do Rio de Janeiro. *Acta Biomedica Brasiliensia*, v.8, n.1, p.35-42, 2017. <DOI: 10.18571/acbm.120>.

- MATESCO, V.C. et al. Contaminação sazonal por ovos de helmintos na praia de Ipanema, em Porto Alegre, Rio Grande do Sul, Brasil. *Revista Agropecuária Tropical*, v.35, n.2, p.135-141, 2006. <DOI: 10.5216/rpt.v35i2.1902>.
- NARASIMHAM, M.V. et al. *Dipylidium caninum* infection in a child: a rare case report. *Indian Journal of Medical Microbiology*, v.31, n.1, p.82-84, 2013. <DOI: 10.4103/0255-0857.108738>.
- NAVONE, G. et al. Estudio transversal de las parasitosis intestinales en poblaciones infantiles de Argentina. *Revista Panamericana de Salud Pública*, v.41, n.2, p.1-9, 2017. <DOI: 10.26633/RPSP.2017.24>.
- OVERGAAUW, P.A.M.; VAN KNAPEN, F. Veterinary and public health aspects of *Toxocara* spp. *Veterinary Parasitology*, v.193, n.4, p.398-403, 2013. <Doi: 10.1016/j.vetpar.2012.12.035>
- PEDRAZA, D.F.; QUEIROZ, D.; SALES, M.C. Doenças infecciosas em crianças pré-escolares brasileiras assistidas em creches. *Ciência e Saúde Coletiva*, v.19, n.2, p.511-528, 2014. <DOI: 0.1590/ 1413-81232014192.09592012>.
- PEREIRA, A. et al. Parasitic zoonoses associated with dogs and cats: a survey of Portuguese pet owners' awareness and deworming practices. *Parasites and Vectors*, v.9, n.1, p.1-9, 2016. <DOI: 10.1186/s13071-016-1533-2>.
- PERUCA, L.C.B.; LANGONI, H.; LUCHEIS, S.B. Larva migrans visceral e cutânea como zoonoses: revisão de leitura. *Veterinária e Zootecnia*, v.16, n.4, p.601-616, 2009.
- PROVIDELO, G.A. et al. Contaminação ambiental por geo-helmintos em praças do município de Botucatu, São Paulo. *Archives of Veterinary Science*, v.25, n.1, p.20-34, 2020. <DOI: 10.5380/avs.v25i1.69181>.
- RIBEIRO, K.L. et al. Avaliação da ocorrência de formas parasitárias no solo de praças públicas do município de Esteio (RS). *Revista Acadêmica: Ciências Agrárias e Ambientais*, v.11, n.1, p.59-64, 2013. <DOI: 10.7213/academica.7756>.
- ROSALES, T.L.L.; MALHEIROS, A.F. Contaminação Ambiental por enteroparasitas presentes em fezes de cães em uma região do Pantanal. *O mundo da saúde*, v.41, n.3, p.368-377, 2017. <DOI: 10.15343/0104-7809.20174103368377>.
- SANTARÉM, V.A.; GIUFFRIDA, R.; ZANIN, G.A. Larva migrans cutânea: ocorrência de casos humanos e identificação de larvas de *Ancylostoma* spp em parque público do município de Taciba, São Paulo. *Revista Brasileira de Medicina Tropical*, v.37, n.2, p.179-181, 2004. <DOI: 10.1590/S0037-86822004000200014>.
- SANTOS, G.M. et al. Investigação soropidemiológica sobre a Larva migrans visceral por *Toxocara canis* em usuários de serviços de saúde de Goiânia-GO. *Revista de Patologia Tropical*, v.38, n.3, p.197-206, 2009. <DOI: 10.5216/rpt.v38i3.7838>.
- SAVIOLI, L.; SMITH, H.; THOMPSON, A. *Giardia* and *Cryptosporidium* join the "Neglected Diseases Initiative". *Trends in Parasitology*, v.22, n.5, p.203-208, 2006. <DOI: 10.1016/j.pt.2006.02.015>.
- SHERDING, R. and JOHNSON, S. Doenças Intestinais. In: BICHARD, S.J. *Manual Saunders de clínica de pequenos animais*. 3 ed. São Paulo: Roca, 2008.
- SILVA, S.M.D.; ARAÚJO, F.A. Prevalência da infecção por *Giardia* sp. em cães do município de Porto Alegre-RS, comparação entre duas populações: cães de rua e cães com proprietário provenientes de áreas de vulnerabilidade social. *Journal of the Health Sciences Institute*, v.31, n.1, p.99-103, 2013.



POTENTIAL ZOO NOTIC PARASITES OF DOGS FECES IN THE VICINITY OF SCHOOLS IN THE CITY OF CAPÃO DO LEÃO, RIO GRANDE DO SUL, BRAZIL / PARASITOS POTENCIALMENTE ZOO NÓTICOS EM FEZES DE CÃES NAS PROXIMIDADES DE ESCOLAS DO MUNICÍPIO DE CAPÃO DO LEÃO, RIO GRANDE DO SUL, BRASIL

SIMONATO, G. et al. Contamination of Italian parks with canine helminth eggs and health risk perception of the public. Preventive Veterinary Medicine. v.172, p. 104788, 2019. <Doi: 10.1016/j. prevetmed.2019.104788>

TRAVERSA, D. et al. Environmental contamination by canine geohelminths. Parasites & Vectors, v.7, n.1, p.67, 2014. <DOI: 10.1186/1756-3305-7-67>.

UCHÔA, C.M.A. et al. Parasitismo intestinal em crianças e funcionários de creches comunitárias na cidade de Niterói-RJ, Brasil. Revista de Patologia Tropical, v.38, n.4, p.267-278, 2009. <DOI: 10.5216/rpt.v38i4.8590>.

WILLIS, I.I. A simple levitation method for the detection of hookworm ova. Medical Journal of Austrália, v.2, n.18, p.375-376, 1921. <DOI:10.5694/j.1326-5377.1921.tb60654.x

ZAGUI, G.S. et al. Adaptação do ecomapa proposto no Modelo Calgary para avaliação socioambiental de parasitoses intestinais em crianças de creches filantrópicas. Revista do Instituto Adolfo Lutz, v.76, p.1-10, 2017.